

# On the universality of the Carter and McLenaghan formula

Ion I. Cotăescu

*West University of Timișoara,  
V. Parvan Ave. 4, RO-300223 Timișoara*

## Abstract

It is shown that the formula of the isometry generators of the spinor representation given by Carter and McLenaghan is universal in the sense that this holds for any representation either in local frames or even in natural ones. The point-dependent spin matrices in natural frames are introduced for any tensor representation deriving the covariant form of the isometry generators in these frames.

Pacs: 04.20.Cv, 04.62.+v, 11.30.-j

## I. INTRODUCTION

In general relativity there are many symmetries but only the isometries (associated to Killing vectors) are able to produce conserved quantities via Noether theorem. The physical fields minimally coupled to gravity take over this symmetry transforming according to appropriate representations of the isometry group. In the case of the scalar vector or tensor fields these representations are completely defined by the well-known rules of the general coordinate transformations since the isometries are in fact particular automorphisms. On the other hand, the theory of spinor fields is formulated in orthogonal local frames where the basis-generators of the spinor representation were discovered by Carter and McLenaghan [1]

Some time ago we proposed a theory of external symmetry in the context of the gauge covariant theories based on local orthogonal frames. We introduced there new transformations which combine isometries and gauge transformations such that the tetrad fields should remain invariant [2]. In this way we obtained the external symmetry group which is nothing else than the universal covering group of the isometry one. Moreover, we derived the general form of the basis-generators of the representations of this group showing that these are given by a similar formula to that of Carter and McLenaghan.

In the present letter we should like to show that this formula holds even in natural frames. To this end we define the spin matrices of the vector and tensor representations in natural frames which are point-dependent and give rise to the spin terms of the covariant form of our basis-generators. The conclusion is that the Carter and McLenaghan formula is universal.

We start in the second section with a short introduction to the tetrad gauge covariant theories in which we defined our theory of external symmetry briefly presented in the next section. The spin matrices in natural frames are introduced in section three where the basis-generators of the vector and tensor representations of the external symmetry group are obtained.

## II. TETRAD GAUGE COVARIANCE

Let us consider the curved spacetime  $M$  and a local chart (natural frame) of coordinates  $x^\mu$  (labeled by natural indices,  $\mu, \nu, \dots = 0, 1, 2, 3$ ). Given a gauge, we denote by  $e_{\hat{\mu}}$  the tetrad fields that define the local frames and by  $\hat{e}^{\hat{\mu}}$  those of the corresponding coframes. These have

the usual orthonormalization properties,  $\hat{e}_{\hat{\alpha}}^{\hat{\mu}} e_{\hat{\nu}}^{\alpha} = \delta_{\hat{\nu}}^{\hat{\mu}}$ ,  $\hat{e}_{\hat{\alpha}}^{\hat{\mu}} e_{\hat{\mu}}^{\beta} = \delta_{\hat{\alpha}}^{\beta}$ ,  $e_{\hat{\mu}} \cdot e_{\hat{\nu}} = \eta_{\hat{\mu}\hat{\nu}}$ ,  $\hat{e}^{\hat{\mu}} \cdot \hat{e}^{\hat{\nu}} = \eta^{\hat{\mu}\hat{\nu}}$ . The metric tensor  $g_{\mu\nu} = \eta_{\hat{\alpha}\hat{\beta}} \hat{e}_{\hat{\mu}}^{\hat{\alpha}} \hat{e}_{\hat{\nu}}^{\hat{\beta}}$  raises or lowers the natural indices while for the local indices ( $\hat{\mu}, \hat{\nu}, \dots = 0, 1, 2, 3$ ) we have to use the Minkowski metric  $\eta = \text{diag}(1, -1, -1, -1)$ . This metric remains invariant under the transformations of its gauge group,  $O(3, 1)$ . This has as subgroup the Lorentz group,  $L_+^{\dagger}$ , of the transformations  $\Lambda[A(\omega)]$  corresponding to the transformations  $A(\omega) \in SL(2, \mathbb{C})$  through the canonical homomorphism. In the standard covariant parametrization, with the real parameters  $\omega^{\hat{\alpha}\hat{\beta}} = -\omega^{\hat{\beta}\hat{\alpha}}$ , the  $SL(2, \mathbb{C})$  transformation read

$$A(\omega) = e^{-\frac{i}{2} \omega^{\hat{\alpha}\hat{\beta}} S_{\hat{\alpha}\hat{\beta}}} \quad (1)$$

where  $S_{\hat{\alpha}\hat{\beta}}$  are the covariant basis-generators of the  $sl(2, \mathbb{C})$  Lie algebra. In fact, these are the principal spin operators which generate the spin terms of other operators in the quantum theory. For small values of  $\omega^{\hat{\alpha}\hat{\beta}}$  the matrix elements of the transformations  $\Lambda$  in the local basis can be expanded as  $\Lambda_{\hat{\nu}}^{\hat{\mu}}[A(\omega)] = \delta_{\hat{\nu}}^{\hat{\mu}} + \omega_{\hat{\nu}}^{\hat{\mu}} + \dots$ .

Assuming now that  $M$  is orientable and time-orientable we can consider  $G(\eta) = L_+^{\dagger}$  as the gauge group of the Minkowski metric  $\eta$  [3]. Then, the fields with spin in local frames transform according to finite-dimensional linear representations,  $\rho$ , of the group  $SL(2, \mathbb{C})$ . In general, a matter field  $\psi_{\rho} : M \rightarrow \mathcal{V}_{\rho}$  is defined over  $M$  with values in the vector space  $\mathcal{V}_{\rho}$  of the representation  $\rho$  which is generally reducible. The covariant derivatives of the field  $\psi_{\rho}$ ,

$$D_{\hat{\alpha}}^{\rho} = e_{\hat{\alpha}}^{\mu} D_{\mu}^{\rho} = \hat{\partial}_{\hat{\alpha}} + \frac{i}{2} \rho(S_{\hat{\gamma}}^{\hat{\beta}\cdot}) \hat{\Gamma}_{\hat{\alpha}\hat{\beta}}^{\hat{\gamma}}, \quad (2)$$

depend on the connection coefficients in local frames,  $\hat{\Gamma}_{\hat{\mu}\hat{\nu}}^{\hat{\sigma}} = e_{\hat{\mu}}^{\alpha} e_{\hat{\nu}}^{\beta} (\hat{e}_{\hat{\gamma}}^{\hat{\sigma}} \Gamma_{\alpha\beta}^{\gamma} - \hat{e}_{\beta,\alpha}^{\hat{\sigma}})$ , which assure the covariance of the whole theory under tetrad gauge transformations. This is the general framework of the theories involving fields with half integer spin which the natural frames are not dealing with.

### III. EXTERNAL SYMMETRIES IN LOCAL FRAMES

A special difficulty in local frames is that the theory is no longer covariant under isometries since these can change the tetrad fields that carry natural indices. For this reason we proposed a theory of external symmetry in which each isometry transformation is coupled to a gauge one able to correct the position of the local frames such that the whole transformation

should preserve not only the metric but the tetrad gauge too [2]. Thus, for any isometry transformation  $x \rightarrow x' = \phi_\xi(x) = x + \xi^a k_a + \dots$ , depending on the parameters  $\xi^a$  ( $a, b, \dots = 1, 2, \dots, N$ ) of the isometry group  $I(M)$ , one must perform the gauge transformation  $A_\xi$  defined as

$$\Lambda_{\hat{\beta}}^{\hat{\alpha}}[A_\xi(x)] = \hat{e}_\mu^{\hat{\alpha}}[\phi_\xi(x)] \frac{\partial \phi_\xi^\mu(x)}{\partial x^\nu} e_\beta^\nu(x) \quad (3)$$

with the supplementary condition  $A_{\xi=0}(x) = 1 \in SL(2, \mathbb{C})$ . Then the transformation laws of our fields are

$$\begin{aligned} e(x) &\rightarrow e'(x') = e[\phi_\xi(x)] \\ (A_\xi, \phi_\xi) : \quad \hat{e}(x) &\rightarrow \hat{e}'(x') = \hat{e}[\phi_\xi(x)] \\ \psi_\rho(x) &\rightarrow \psi'_\rho(x') = \rho[A_\xi(x)] \psi_\rho(x). \end{aligned} \quad (4)$$

We have shown that the pairs  $(A_\xi, \phi_\xi)$  constitute a well-defined Lie group we called the external symmetry group,  $S(M)$ , pointing out that this is just the universal covering group of  $I(M)$  [2]. For small values of  $\xi^a$ , the  $SL(2, \mathbb{C})$  parameters of  $A_\xi(x) \equiv A[\omega_\xi(x)]$  can be expanded as  $\omega_\xi^{\hat{\alpha}\hat{\beta}}(x) = \xi^a \Omega_a^{\hat{\alpha}\hat{\beta}}(x) + \dots$ , in terms of the functions

$$\Omega_a^{\hat{\alpha}\hat{\beta}} \equiv \frac{\partial \omega_\xi^{\hat{\alpha}\hat{\beta}}}{\partial \xi^a} \Big|_{\xi=0} = (\hat{e}_\mu^{\hat{\alpha}} k_{a,\nu}^\mu + \hat{e}_{\nu,\mu}^{\hat{\alpha}} k_a^\mu) e_\lambda^\nu \eta^{\lambda\hat{\beta}} \quad (5)$$

which depend on the Killing vectors  $k_a = \partial_{\xi^a} \phi_\xi|_{\xi=0}$  associated to  $\xi^a$ .

The last of Eqs. (4) defines the operator-valued representation  $(A_\xi, \phi_\xi) \rightarrow T_\xi^\rho$  of the group  $S(M)$  whose transformations,  $(T_\xi^\rho \psi_\rho)[\phi_\xi(x)] = \rho[A_\xi(x)] \psi_\rho(x)$ , leave the field equation invariant. Their basis-generators [2],

$$X_a^\rho = i \partial_{\xi^a} T_\xi^\rho \Big|_{\xi=0} = -i k_a^\mu \partial_\mu + \frac{1}{2} \Omega_a^{\hat{\alpha}\hat{\beta}} \rho(S_{\hat{\alpha}\hat{\beta}}), \quad (6)$$

commute with the operator of the field equation and satisfy the commutation rules  $[X_a^\rho, X_b^\rho] = i c_{abc} X_c^\rho$  determined by the structure constants,  $c_{abc}$ , of the  $s(M) \sim i(M)$  algebras. Eq. (6) can be put in covariant the form

$$X_a^\rho = -i k_a^\mu D_\mu^\rho + \frac{1}{2} k_{a\mu;\nu} e_\alpha^\mu e_\beta^\nu \rho(S^{\hat{\alpha}\hat{\beta}}), \quad (7)$$

which represents a generalization to any representation  $\rho$  of the famous formula given by Carter and McLenaghan for the spinor representation  $\rho_{spin} = (\frac{1}{2}, 0) \oplus (0, \frac{1}{2})$  [1].

#### IV. ISOMETRIES IN NATURAL FRAMES

Whenever the spinors are absent, the matter fields are vectors or tensors of different ranks and the whole theory is independent on the tetrad fields dealing with the natural frames only. Let us see what happens with our theory of external symmetry in natural frames focussing on tensor fields. Any tensor field,  $\Theta$ , transforms under isometries as  $\Theta \rightarrow \Theta' = T_\xi \Theta$ , according to a tensor representation of the group  $S(M)$  defined by the well-known rule in natural frames

$$\left[ \frac{\partial \phi_\xi^\alpha(x)}{\partial x_\mu} \frac{\partial \phi_\xi^\beta(x)}{\partial x_\nu} \dots \right] (T_\xi \Theta)_{\alpha\beta\dots} [\phi(x)] = \Theta_{\mu\nu\dots}(x). \quad (8)$$

Hereby one derives the basis-generators of the tensor representation,  $X_a = i \partial_{\xi_a} T_\xi|_{\xi=0}$ , whose action reads

$$(X_a \Theta)_{\alpha\beta\dots} = -i(k_a^\nu \Theta_{\alpha\beta\dots;\nu} + k_a^\nu{}_{;\alpha} \Theta_{\nu\beta\dots} + k_a^\nu{}_{;\beta} \Theta_{\alpha\nu\dots\dots}). \quad (9)$$

Our purpose is to show that the operators  $X_a$  can be written in a form which is equivalent to Eq. (7) of Carter and McLanaghan.

We start with the vector representation  $\rho_v = (\frac{1}{2}, \frac{1}{2})$  of the  $SL(2, \mathbb{C})$  group, generated by the spin matrices  $\rho_v(S^{\hat{\alpha}\hat{\beta}})$  which have the well-known matrix elements

$$[\rho_v(S^{\hat{\alpha}\hat{\beta}})]_{\hat{\nu}}^{\hat{\mu}} = i(\eta^{\hat{\alpha}\hat{\mu}} \delta_{\hat{\nu}}^{\hat{\beta}} - \eta^{\hat{\beta}\hat{\mu}} \delta_{\hat{\nu}}^{\hat{\alpha}}), \quad (10)$$

in local bases. Furthermore, we define the point-dependent spin matrices in natural frames whose matrix elements in the natural basis read

$$(\tilde{S}^{\mu\nu})_{\tau}^{\sigma} = e_{\hat{\alpha}}^{\mu} e_{\hat{\beta}}^{\nu} e_{\hat{\gamma}}^{\sigma} [\rho_v(S^{\hat{\alpha}\hat{\beta}})]_{\hat{\delta}}^{\hat{\gamma}} \hat{e}_{\tau}^{\hat{\delta}} = i(g^{\mu\sigma} \delta_{\tau}^{\nu} - g^{\nu\sigma} \delta_{\tau}^{\mu}). \quad (11)$$

In what follows we assume that these matrices represent the spin operators of the vector representation in natural frames. We observe that these are the basis-generators of the groups  $G[g(x)] \sim G(\eta)$  which leave the metric tensor  $g(x)$  invariant in each point  $x$ . Since the representations of these groups are point-wise equivalent with those of  $G(\eta)$ , one can show that in each point  $x$  the basis-generators  $\tilde{S}^{\mu\nu}(x)$  satisfy the standard commutation rules of the vector representation  $\rho_v$  (but with  $g(x)$  instead of  $\eta$ ).

In general, the spin matrices of a tensor  $\Theta$  of any rank,  $n$ , are the basis-generators of the representation  $\rho_n = \rho_v^1 \otimes \rho_v^2 \otimes \rho_v^3 \otimes \dots \otimes \rho_v^n$  which read  $\rho_n(\tilde{S}) = \tilde{S}^1 \otimes I^2 \otimes I^3 \dots + I^1 \otimes \tilde{S}^2 \otimes I^3 \dots + \dots$ . Using these spin matrices a straightforward calculation shows that Eq. (7) can be rewritten

in natural frames as

$$X_a^n = -ik_a^\mu \nabla_\mu + \frac{1}{2} k_{a\mu;\nu} \rho_n(\tilde{S}^{\mu\nu}), \quad (12)$$

where  $\nabla_\mu$  are the usual covariant derivatives. It is not difficult to verify that the action of these operators is just that given by Eq. (9) which means that  $X_a^n$  are the basis-generators of a tensor representation of rank  $n$  of the group  $S(M)$ . Thus, it is clear that Eq. (12) represents the generalization to natural frames of the Carter and McLenaghan formula (7).

## V. CONCLUDING REMARKS

The principal conclusion here is that the Carter and McLenaghan formula is universal since it holds not only in local frames but in natural frames too. In local frames this formula can be written in the covariant form (7) or as in Eq. (6) where the orbital part (i.e. the first term) is completely separated from the spin term (the second one). It is worth pointing out that Eq. (6) can not be rewritten in natural frames even though it might be useful for analyzing the operators structure in the relativistic quantum mechanics. This is because the connection in local frames can not be expressed in terms of spin matrices  $\tilde{S}$ . Therefore, in natural frames we are forced to use only the genuine Carter and McLenaghan formula (12).

This formula exhibits explicitly the point-dependent spin matrices of the vector or tensor representations which may provide us with information about the spin content of the various fields involved in theory. However, some results could be surprising. Thus, for example, if a second order rank tensor has the form  $G_{\mu\nu} = g_{\mu\nu}F$  where  $F$  is a scalar field then we calculate  $\rho_2(\tilde{S}^{\alpha\beta})G = 0$  which indicates that the tensor  $G$  is spinless despite of the fact that it is of second rank. Consequently, the metric tensor is also spinless as it seems to be natural since this is covariantly constant and invariant under isometries (such that  $X_a^2 g = 0$ ). Another example is presented in Ref. [4] where we define the photon polarization in de Sitter manifolds without to consider local frames.

Finally we note that, despite of our partial results, the problem of the spin in general relativity remains open as long as we do not have an effective theory similar to the Wigner theory of the induced representations of the Poincaré group.

## Acknowledgments

We are grateful to Cosmin Crucean and Nistor Nicolaevici for interesting and useful discussions on closely related subjects.

---

- [1] B. Carter and R. G. McLenaghan, *Phys. Rev. D* **19**, 1093 (1979).
- [2] I. I. Cotăescu, *J. Phys. A: Math. Gen.* **33**, 9177 (2000).
- [3] R. M. Wald, *General Relativity*, (Univ. of Chicago Press, Chicago and London, 1984)
- [4] I. I. Cotăescu and C. Crucean, [arXiv:0806.2515](#) [gr-qc]